arXiv:1303.6002v2 [math.AG] 28 Mar 2013

UNIFORMLY NONSPECIAL LINEAR SYSTEMS ON NODAL CURVES

ZIV RAN

Abstract. A line bundle having large degree on every subcurve of a given nodal curve is well-behaved.

On a smooth (more generally, irreducible nodal) curve X of genus g, it is trivial or well known [3], [4] that a line bundle L of degree d is well-behaved when d is large enough relative to g:

(i) if $d \geq 2g-1$, then $h^0(L) = d - g + 1$;
(ii) if $d \geq 2g$, L is globally generated;
(iii) if $d \geq 2g+1$, then L is very ample.
(iv) if $d \geq 2g+1+p$, then L has property N_p (theorem of Mrk Green, see [2] or [4], §1.8).

The obvious generalizations of these properties to the case of nodal, possibly reducible, curves trivially fails. Clearly, any numerical condition guaranteeing good behavior on reducible curves must involve subcurves of X. One such condition called *balancedness* was considered by Caporaso [1]. In this paper we focus on a different numerical condition that we call *uniform numerical nonspeciality*. In general, this notion is neither stronger nor weaker than balancedness, but for large degree it is weaker. Based on the degree of uniform nonspeciality, we prove a clean generalization of properties (i-iv) above. As a (very special) consequence, we obtain a short proof of one of the main results of [1].

In this paper, all curves are nodal, possibly disconnected. For a nodal curve X, the genus $g(X)$ is the arithmetic genus, i.e. the number such that $\deg(\omega_X) = 2g(X) - 2$.

Definition 1. *(i) A line bundle L on X is said to be k- numerically nonspecial or k-nuns if*

$$\deg(L) \geq 2g(X) - 2 + k.$$

(ii) L is said to be k-uniformly numerically nonspecial or k-ununs if for all subcurves $Y \subset X$, L_Y is k-nuns on Y.

(iii) L is said to be k-spanned if for every ideal I of colength k on X, the natural map $H^0(L) \to L/IL$ is surjective. □

Date: April 1, 2013.
1991 *Mathematics Subject Classification.* 14N99,14H99.
Key words and phrases. linear systems, nodal curves.

Thus, 1-spanned is globally generated, 2-spanned is very ample, etc. It is well known that for X irreducible nodal, k-nuns implies $(k-1)$-spanned. Our main result here is that for all X nodal, k-ununs implies $(k-1)$-spanned and N_{k-3}.

Lemma 2. *Let p be a node on X, let $J_{a,b}$ be the ideal of type (x^a, y^b) of colength $e = a+b-1$ cosupported at p, let $\pi : X' \to X$ be the blowing-up of p, and L' the unique line bundle on X' of degree $\deg(L) - a - b$ such that $\pi_*(L') = J_{a,b}L$. If L is k-ununs on X, then L' is $(k-e)$-ununs on X'.*

Proof. We may assume $0 < b \le a \le e$. Let Y' be a subcurve of X' and suppose to begin with that Y' contains both preimages of p. Then Y' is the blowing-up of p on a uniquely determined subcurve Y of X containing p, and we have

$$\deg(L'_{Y'}) = \deg(L_Y) - a - b \ge 2g(Y) - 2 + k - a - b = 2g(Y') - 2 + k + 1 - e,$$

which is good enough. If Y' contains precisely one preimage of p, it maps isomorphically to a subcurve Y of X through p and we have

$$\deg(L'_{Y'}) \ge d - a \ge 2g(Y) - 2 + k - a \ge 2g(Y') - 2 + k - e$$

as $a \le e$. If Y' contains no preimage of p then $L|_{Y'}$ is trivially k-nuns. This concludes the proof. □

Theorem 3. *(i) If L is 1-ununs , then $h^0(L) = \deg(L) - g + 1$.*
(ii) If L is k-ununs for $k > 1$, then L is $(k-1)$-spanned.
(iii) If L is k-ununs, $k \ge 3$, then L has property N_{k-3}.

Proof. (i) By Riemann-Roch and Serre Duality, it suffices to prove that $h^0(\omega_X(-L)) = 0$. This is proved by induction on the number of irreducible components of X. If X is irreducible, the result is clear. For the induction step, let $s \in H^0(\omega_X(-L))$. Note that

$$\deg(L) = \sum_Z \deg(L_Z) > \deg(\omega_X) = \sum_Z \deg(\omega_X|_Z)$$

the sum being over all irreducible components Z of X. Therefore there exists an irreducible component Z such that

$$\deg(L_Z) > \deg(\omega_X|_Z) \ge \deg(\omega_Z).$$

Therefore $s|_Z = 0$. Let $Y = \overline{X \setminus Z}$ be the complementary curve. Then since s vanishes on $Y \cap Z$, it may be viewed as a section of $\omega_Y(-L)$. Since $L|_Y$ is 1-ununs, it follows that $s = 0$.

(ii) We need to show that for any ideal J of colength $e < k$ on X, $h^0(JL) = \deg(L) - g(X) + 1 - e$. If J is invertible, we just apply Part (i) to JL. Therefore we may assume J is cosupported at the nodes. Using an obvious induction, we may assume J is cosupported at a single node $p \in X$. If J is invertible, again we may conclude by applying Part (i) to JL. If not, the results of [5] show that J has the form $J_{a,b}$ as in Lemma 2 and it suffices to apply Part (i) to L' on X'.

(iii) Given that L is very ample and satisfies (i), the proof of Theorem (4.a.1) in [2] goes through essentially verbatim. More specifically, property N_{k-3} is equivalent to the vanishing of the Koszul cogomology $K_{k-3,2}(X, L)$. By an evident extension of Green's Duality Theorem (2.c.6), the latter group is dual to $K_{h^0(L)+1-k,0}(X, \omega_X, L)$, which vanishes by (an evident extension of) Green's vanishing theorem (3.a.1). $\square$

The above Theorem- in fact, just the 1-ununs and 2-ununs cases- yield a quick proof and strengthening of Caporaso's Theorem 2.2.1 in [1], which concerns *balanced* line bundles. A line bundle L of degree d on a semistable curve X is said to be balanced if for every subcurve $Z \subset X$ meeting its complementary curve in δ_Z points, we have

$$d\frac{2g(Z)-2+\delta_Z}{2g(X)-2} - \delta_Z/2 \leq \deg(L_Z) \leq d\frac{2g(Z)-2+\delta_Z}{2g(X)-2} + \delta_Z/2,$$

and moreover L has positive degree on each nonsingular rational component containing 2 nodes of X. If L is balanced of nonnegative degree, then it has nonnegative degree on every subcurve. Now suppose L is balanced of degree $d > 2g(X) - 2$. Then in the above inequality we have

$$\deg(L_Z) > 2g(Z) - 2 + \delta_Z - \delta_Z/2 = 2g(Z) - 2 + \delta_Z/2.$$

It follows that L is 1-ununs, hence $h^0(L) = d - g(X) + 1$. This is Theorem 2.2.1 (i) of [1]. Now suppose moreover that X has no separating nodes and that $\deg(L) \geq 2g(X)$. Then for every proper subcurve $Z \subsetneq X$, we have $\delta_Z \geq 2$, hence $\deg(L_Z) \geq 2g(Z)$. If moreover $d \geq 2g(X)$, it follows that L is 2-ununs, hence by Part (ii) above is globally generated. This is Theorem 2.2.1(ii) of [1]. Finally, assume X is semistable of genus 2 or more, and L is balanced of degree $d \geq 5(g(X) - 1)$. Let $Z \subset X$ be a subcurve. We claim that $\deg(L_Z) \geq 2g(Z)$. To prove this, it is easy to see that no generality is lost by assuming X, Z are connected, $g(Z) > 0$ and $Z \subsetneq X$. By balancedness we have

$$\deg(L_Z) \geq 2g(Z) - 2 + \frac{3}{2}(2g(Z) - 2) + 2\delta_Z.$$

If $g(Z) > 0$, then $\delta_Z > 0$ and the last two terms add up to at least 2. This proves our claim. Hence L is 2-ununs and therefore globally generated. This is Theorem 2.2.1(iii) of [1].

REFERENCES

1. L. Caporaso, *Linear series on semistable curves*, International math. res. notices **2011**, 1107–1134, `arxiv.org:0812.1682v4`.
2. M. L. Green, *Koszul cohomology and the geometry of projective varieties*, J. Differential Geometry **19** (1984), 125–171.
3. R. Hartshorne, *Algebraic Geometry*, Springer, Berlin, 1977.
4. R. Lazarsfeld, *Positivity in algebraic geometry*, Springer, 2004.
5. Z. Ran, *A note on Hilbert schemes of nodal curves*, J. Algebra **292** (2005), 429–446.

UC MATH DEPT.
BIG SPRINGS ROAD SURGE FACILITY
RIVERSIDE CA 92521

E-mail address: `ziv.ran @ucr.edu`